

## Statistics Formula Sheet:

<b>Mean:</b>	<b>Sample Mean:</b> $\bar{X} = \frac{\sum X}{n}$	<b>Population Mean:</b> $u = \frac{\sum X}{N}$
<b>Median: (Q2)</b>	<b>If n is odd:</b> $M = \left(\frac{n+1}{2}\right)^{th} \text{ Term}$	<b>If n is even:</b> $M = \frac{\left(\frac{n}{2}\right)^{th} \text{ Term} + \left(\frac{n}{2} + 1\right)^{th} \text{ Term}}{2}$
<b>Mode:</b>	The number with the highest frequency.	
<b>Range:</b> H → Highest Value L → Lowest Value	$\text{Range} = H - L$	$\text{MidRange} = \frac{H + L}{2}$
<b>Standard Deviation:</b>	<b>Sample:</b> $s = \sqrt{\frac{\sum(X - \bar{X})^2}{n - 1}}$	<b>Population:</b> $\sigma = \sqrt{\frac{\sum(X - u)^2}{N}}$
<b>Variance:</b>	<b>Sample Variance:</b> $s^2 = \frac{\sum(X - \bar{X})^2}{n - 1}$	<b>Population Variance:</b> $\sigma^2 = \frac{\sum(X - u)^2}{N}$
<b>Coefficient of Variation:</b>	<b>Sample CV:</b> $CV = \frac{s}{\bar{X}} \times 100\%$	<b>Population CV:</b> $CV = \frac{\sigma}{u} \times 100\%$
<b>Mean Absolute Deviation:</b>	<b>Sample MD (Mean):</b> $MD = \frac{\sum X - \bar{X} }{n}$	<b>Population MD (Mean):</b> $MD = \frac{\sum X - u }{N}$
<b>Average Deviation:</b>	<b>Sample AD:</b> $AD = \frac{\sum(X - \bar{X})}{n}$	<b>Population AD:</b> $AD = \frac{\sum(X - u)}{N}$

<b>Quartile:</b>	$Q_k = k \left( \frac{n+1}{4} \right)^{th} \text{ Term}$ $Q_1 = 1 \left( \frac{n+1}{4} \right)^{th} \text{ Term} \quad Q_3 = 3 \left( \frac{n+1}{4} \right)^{th} \text{ Term}$
<b>Percentile:</b>	$P_k = k \left( \frac{n+1}{100} \right)^{th} \text{ Term}$ $P_{30} = 30 \left( \frac{n+1}{100} \right)^{th} \text{ Term} \quad P_{70} = 70 \left( \frac{n+1}{100} \right)^{th} \text{ Term}$
<b>Decile:</b>	$D_k = k \left( \frac{n+1}{10} \right)^{th} \text{ Term}$
<b>Octile:</b>	$O_k = k \left( \frac{n+1}{8} \right)^{th} \text{ Term}$
<b>Interquartile Range:</b>	$IQR = Q_3 - Q_1$
<b>Quartile Deviation:</b>	$QD = \frac{Q_3 - Q_1}{2} = \frac{1}{2}(IQR)$
<b>Coefficient of Quartile Deviation:</b>	$CQD = \frac{Q_3 - Q_1}{Q_3 + Q_1}$
<b>Range of Outliers:</b>	$[Q_1 - 1.5 IQR, Q_3 + 1.5 IQR]$ <p><b>Note:</b> Any data point that exists outside of the range shown above is considered an outlier.</p>
<b>Coefficient of Range:</b>	$CR = \frac{H - L}{H + L}$

	General Formula:	Expanded Form:	2 Numbers:
<b>Arithmetic Mean:</b>	$\bar{X} = \frac{\sum X}{n}$	$\bar{X} = \frac{X_1 + X_2 + X_3 + \dots + X_n}{n}$	$AM = \frac{a + b}{2}$
<b>Geometric Mean:</b>	$\bar{X}_G = \left( \prod_{i=1}^n X_i \right)^{\frac{1}{n}}$	$\bar{X}_G = (X_1 * X_2 * X_3 \dots X_n)^{1/n}$	$GM = \sqrt{ab}$
	$\bar{X}_G = 10^{\left( \frac{\sum \log(X)}{n} \right)}$	$\bar{X}_G = 10^{\left( \frac{\log(X_1) + \log(X_2) + \dots + \log(X_n)}{n} \right)}$	$GM = 10^{\frac{\log(a) + \log(b)}{2}}$
<b>Weighted Mean:</b>	$\bar{X}_W = \frac{\sum WX}{W}$	$\bar{X}_W = \frac{W_1X_1 + W_2X_2 + \dots + W_nX_n}{W_1 + W_2 + \dots + W_n}$	$WM = \frac{W_1a + W_2b}{W_1 + W_2}$
<b>Harmonic Mean:</b>	$\bar{X}_H = \frac{n}{\sum \left( \frac{1}{X} \right)}$	$\bar{X}_H = \frac{n}{\frac{1}{X_1} + \frac{1}{X_2} + \frac{1}{X_3} + \dots + \frac{1}{X_n}}$	$HM = \frac{2}{\frac{1}{a} + \frac{1}{b}} = \frac{2ab}{a + b}$
<b>Root Mean Square:</b>	$X_{rms} = \sqrt{\frac{\sum (X^2)}{n}}$	$X_{rms} = \sqrt{\frac{X_1^2 + X_2^2 + X_3^2 + \dots + X_n^2}{n}}$	$X_{rms} = \sqrt{\frac{a^2 + b^2}{2}}$
<b>Mean Relationship:</b>	$GM = \sqrt{(AM)(HM)} \quad \text{For 2 Numbers}$ $\sqrt{ab} = \sqrt{\left( \frac{a+b}{2} \right) \left( \frac{2ab}{a+b} \right)}$		

## Statistics Formulas for Grouped Data:

<b>Mean:</b>	$\bar{X} = \frac{\sum fX_m}{\sum f} = \frac{\sum fX_m}{n}$
<b>Midpoint of Range:</b>	$X_m = \frac{X_1 + X_2}{2}$
<b>Standard Deviation:</b>	$s = \sqrt{\frac{\sum f(X_m - \bar{X})^2}{n - 1}} = \sqrt{\frac{\sum fX_m^2 - \frac{(\sum fX_m)^2}{n}}{n - 1}}$
<b>Variance:</b>	$s^2 = \frac{\sum f(X_m - \bar{X})^2}{n - 1} = \frac{\sum fX_m^2 - \frac{(\sum fX_m)^2}{n}}{n - 1}$
<b>1<sup>st</sup> Quartile:</b>	$Q_1 = L_1 + \frac{w_1}{f_1} \left( \frac{n}{4} - C_1 \right)$ <p style="text-align: center; margin-top: 5px;"> <i>L</i> → Lower Class Boundary      <i>w</i> → Width of Class Interval         </p>
<b>Median – 2<sup>nd</sup> Quartile:</b>	$\text{Median} = Q_2 = L_2 + \frac{w_2}{f_2} \left( \frac{n}{2} - C_2 \right)$ <p style="text-align: center; margin-top: 5px;"> <i>f</i> → frequency of quartile class      <i>n</i> → total frequency         </p>
<b>3<sup>rd</sup> Quartile:</b>	$Q_3 = L_3 + \frac{w_3}{f_3} \left( \frac{3n}{4} - C_3 \right)$ <p style="text-align: center; margin-top: 5px;"> <i>C</i> → Cumulative frequency of preceding quartile class.         </p>
<b>Mode:</b>	$\text{Mode} = L + h \left( \frac{f_m - f_1}{2f_m - f_1 - f_2} \right)$ <p style="margin-top: 5px;"> <i>L</i> → Lower boundary of modal class  <i>h</i> → Size of class interval  <i>f<sub>m</sub></i> → frequency of modal class  <i>f<sub>1</sub></i> → frequency of preceding class  <i>f<sub>2</sub></i> → frequency of succeeding class         </p>